



Point Line Configurations and their Realizability

Brigitte Servatius



Home Page

Print

Title Page



Page 1 of 36

Go Back

Full Screen

Close

Quit



Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page

◀◀ ▶▶

◀ ▶

Page 2 of 36

Go Back

Full Screen

Close

Quit

Zur Theorie der Netze und Configurationen von Konrad Zindler in Graz

- Elementary proof of a theorem of Möbius:

Given 4 points in the plane, one can, by ruler alone construct a point in the ϵ -neighborhood of a given 5'th point for any $\epsilon > 0$.

- Generalization of Configuration:

A system of points and lines in the plane such that on every line there are at least 3 points and through every point there are at least 3 lines.



Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page



Page 3 of 36

Go Back

Full Screen

Close

Quit

1. Zindler's Construction



Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page



Page 4 of 36

Go Back

Full Screen

Close

Quit





Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page





Page 5 of 36

Go Back

Full Screen

Close

Quit





Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page




Page 6 of 36

Go Back

Full Screen

Close

Quit





Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page




Page 7 of 36

Go Back

Full Screen

Close

Quit





Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page



Page 8 of 36

Go Back

Full Screen

Close

Quit





Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page




Page 9 of 36

Go Back

Full Screen

Close

Quit



Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page




Page 10 of 36

Go Back

Full Screen

Close

Quit





Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page




Page 11 of 36

Go Back

Full Screen

Close

Quit





Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page




Page 12 of 36

Go Back

Full Screen

Close

Quit





Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page




Page 13 of 36

Go Back

Full Screen

Close

Quit





Realizable Moves

Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page



Page 14 of 36


Go Back

Full Screen


Close

Quit


- Put a new point on a line.




- Put a new line through a point.



- Intersect two lines.



- Draw a line through two points.




- Join two components by putting a point of one component on a line of the other component.



Realizable Moves

- Put a new point on a line.
- Put a new line through a point.
- Intersect two lines.
- Draw a line through two points.
- Join two components by putting a point of one component on a line of the other component.



Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page



Page 15 of 36

Go Back

Full Screen

Close

Quit



Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page



Page 16 of 36

Go Back


Full Screen

Close


Quit

Dangerous Moves

- Intersect two lines.



- Draw a line through two points.





Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page



Page 17 of 36

Go Back

Full Screen

Close


Quit

Realizable Moves on the Levi graph

- Add vertices of degree one.
- Add vertices of degree two such that bipartiteness and girth 6 are preserved.
(between points of the same color a distance at least 4 apart.)
- Add edges between connected components (bridges).



Given a bipartite graph G of girth 6, these moves may be reversed, provided there exists a vertex of degree at most 2. If G is 3-regular, then $G - v$ can be built up from a vertex by allowable moves



Zindler's ...
Whiteley's Theorem

Home Page

Print

Title Page

◀ ▶

◀ ▶

Page 18 of 36

Go Back


Full Screen

Close

Quit




Given a bipartite graph G of girth 6, these moves may be reversed, provided there exists a vertex of degree at most 2. If G is 3-regular, then $G - v$ can be built up from a vertex by allowable moves





Given a bipartite graph G of girth 6, these moves may be reversed, provided there exists a vertex of degree at most 2. If G is 3-regular, then $G - v$ can be built up from a vertex by allowable moves



Zindler's ...
Whiteley's Theorem

Home Page

Print

Title Page

◀ ▶

◀ ▶

Page 20 of 36

Go Back


Full Screen

Close

Quit



Given a bipartite graph G of girth 6, these moves may be reversed, provided there exists a vertex of degree at most 2. If G is 3-regular, then $G - v$ can be built up from a vertex by allowable moves



Zindler's ...
Whiteley's Theorem

Home Page

Print

Title Page

◀ ▶

◀ ▶

Page 21 of 36

Go Back


Full Screen

Close

Quit



Given a bipartite graph G of girth 6, these moves may be reversed, provided there exists a vertex of degree at most 2. If G is 3-regular, then $G - v$ can be built up from a vertex by allowable moves



Zindler's ...
Whiteley's Theorem

Home Page

Print

Title Page

◀ ▶

◀ ▶

Page 22 of 36

Go Back


Full Screen

Close

Quit



Given a bipartite graph G of girth 6, these moves may be reversed, provided there exists a vertex of degree at most 2. If G is 3-regular, then $G - v$ can be built up from a vertex by allowable moves



Zindler's ...
Whiteley's Theorem

Home Page

Print

Title Page

◀ ▶

◀ ▶

Page 23 of 36

Go Back


Full Screen

Close

Quit



Given a bipartite graph G of girth 6, these moves may be reversed, provided there exists a vertex of degree at most 2. If G is 3-regular, then $G - v$ can be built up from a vertex by allowable moves



Zindler's ...
Whiteley's Theorem

Home Page

Print

Title Page

◀ ▶

◀ ▶

Page 24 of 36

Go Back


Full Screen

Close

Quit



Given a bipartite graph G of girth 6, these moves may be reversed, provided there exists a vertex of degree at most 2. If G is 3-regular, then $G - v$ can be built up from a vertex by allowable moves



Zindler's ...
Whiteley's Theorem

Home Page

Print

Title Page

◀ ▶

◀ ▶

Page 25 of 36

Go Back

Full Screen

Close

Quit



Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page




Page 26 of 36

Go Back

Full Screen

Close

Quit





Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page




Page 27 of 36

Go Back

Full Screen

Close

Quit





Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page




Page 28 of 36

Go Back

Full Screen

Close

Quit





Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page



Page 29 of 36

Go Back

Full Screen

Close

Quit

Steinitz's Theorem

Every symmetric v_3 configuration has a realization in the plane with at most one curved line.



Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page




Page 30 of 36

Go Back

Full Screen

Close

Quit





Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page



Page 31 of 36

Go Back

Full Screen

Close

Quit

Grünbaum's Conjecture

Steinitz's Theorem is true for configurations whose Levi graph is 3-connected.

Theorem

Steinitz's Theorem is true for configurations whose Levi graph is 3-connected. and edge 4-connected.




2. Whiteley's Theorem

A generic picture in $k - 1$ space of an incidence structure lifts to a sharp scene in k -space if and only if

$$i \leq a + kb - (k + 1)$$

for all sub-incidence structures having at least two blocks.



Zindler's ...
Whiteley's Theorem

Home Page

Print

Title Page

◀ ▶

◀ ▶

Page 32 of 36

Go Back

Full Screen

Close

Quit



For a 3-regular bipartite graph of girth six Whiteley's count is violated by three.

Zindler's ...
Whiteley's Theorem

Home Page

Print

Title Page




Page 33 of 36

Go Back

Full Screen

Close

Quit





For a 3-regular bipartite graph of girth six Whiteley's count is violated by three.

Zindler's ...

Whiteley's Theorem

Home Page

Print

Title Page




Page 34 of 36

Go Back


Full Screen

Close

Quit



$i = 6$	$p = 2$	$l = 6$	$l + 2p - 2 = 8$	$2l + p - 2 = 12$
$i = 18$	$p = 8$	$l = 6$	$l + 2p - 2 = 20$	$2l + p - 2 = 18$
$i = 24$	$p = 8$	$l = 9$	$l + 2p - 2 = 23$	$2l + p - 2 = 24$
$i = 27$	$p = 9$	$l = 9$	$l + 2p - 2 = 25$	$2l + p - 2 = 25$




$i = 3$	$p = 1$	$l = 6$	$l + 2p - 2 = 8$	$2l + p - 2 = 12$
$i = 9$	$p = 8$	$l = 6$	$l + 2p - 2 = 20$	$2l + p - 2 = 18$
$i = 21$	$p = 8$	$l = 9$	$l + 2p - 2 = 23$	$2l + p - 2 = 24$
$i = 27$	$p = 9$	$l = 9$	$l + 2p - 2 = 25$	$2l + p - 2 = 25$
$i = 30$	$p = 9$	$l = 9$	$l + 2p - 2 = 25$	$2l + p - 2 = 25$



An (8_4) spatial configuration.
 $a = 8, b = 8, i = 32,$

$$a + 3b - 4 = 28$$



Zindler's ...
Whiteley's Theorem

Home Page

Print

Title Page

Navigation arrows

Navigation arrows

Page 35 of 36

Go Back

Full Screen

Close

Quit



A similar (8_4) spatial configuration.
 Levi graph is a hypercube
 $a = 8, b = 8, i = 32,$

$$a + 3b - 4 = 28$$

Zindler's ...
 Whiteley's Theorem

Home Page

Print

Title Page

◀ ▶

◀ ▶

Page 36 of 36

Go Back

Full Screen

Close

Quit

