

Combinatorial Maps

- The flag graph
- Vertex splitting . . .
- Reduction to . . .
- Classification of . .
- Joining two maps
- Regular maps on . . .
- The 3-block tree
- Matroids
- $\Delta$ -matroids



Full Screen

Close

Quit

### Combinatorial Maps and delta-matroids

Remi Cocou Avohou, Brigitte Servatius and Herman Servatius Worcester Polytechnic Institute



Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Classification of... Classification of... Classification of... Classification of... The 3-block tree Matroids A-matroids

Home Page	
Title Page	
<b>44 &gt;&gt;</b>	
◀ ▶	
Page 2 of <mark>36</mark>	
Go Back	

Close

Quit

Full Screen

# 1. Combinatorial Maps

Tutte [8] defined maps axiomatically. If we have three fixed point free permutations  $\tau_0$ ,  $\tau_2$ , and  $\mathcal{V}$  on a set  $\Phi$  of *flags* such that

- A1  $\tau_0^2 = \tau_2^2 = \operatorname{Id}$
- $\mathbf{A2} \ \tau_0 \tau_2 = \tau_2 \tau_0$
- A3  $\mathcal{V} au_2 = au_2 \mathcal{V}^{-1}$
- ${f A4}\,\left\{{\cal V}^i\phi
  ight\}\cap\left\{{\cal V}^i au_2\phi
  ight\}=\emptyset$

A5  $\tau_0$ ,  $\tau_2$  and  $\tau_0 \tau_2$  are fixed point free

A6  $\langle \tau_0, \tau_2, \mathcal{V} \rangle$  acts transitively on  $\Phi$ 

then we can define a graph G whose vertices are the orbits of  $\Phi$ under  $\langle \tau_2, \mathcal{V} \rangle$  and whose edges are the orbits of  $\Phi$  under  $\langle \tau_0, \tau_2 \rangle$ . The orbits of  $\langle \tau_0, \tau_2 \rangle$  each have four elements and intersect either one or two orbits of  $\langle \tau_2, \mathcal{V} \rangle$ , defining the endpoints of the edge.  $M(G, \tau_0, \tau_2, \mathcal{V})$  is a combinatorial map.



#### Combinatorial Maps

- The flag graph
- Vertex splitting ...
- Reduction to . . .
- Classification of . . .
- Joining two maps
- Regular maps on . . .
- The 3-block tree
- Matroids
- $\Delta$ -matroids







Page <mark>3</mark> of <mark>36</mark>



Full Screen

Close

Quit



Decomposition of the hexagonal prism into flags.



- Combinatorial Maps
- The flag graph
- Vertex splitting ...
- Reduction to . . .
- Classification of . . .
- Joining two maps
- Regular maps on . . .
- The 3-block tree
- Matroids
- $\Delta$ -matroids
  - Home Page





Go Back

Full Screen

Close

Quit



Orbits under  $\tau_2 \tau_0$ ,  $\tau_2 \tau_1$  and  $\tau_0 \tau_1$ .



- Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps Regular maps on... The 3-block tree Matroids Δ-matroids
- Home Page
  Title Page





Full Screen

Close

Quit





Flags at an edge, and at a vertex.



#### Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps Regular maps on...

The	3-h	lock	tree
	U 2.		

Matroids

1	_	n	7.	а	t	r	0	ic	1	s	

Home Page	
Title Page	

••	••
•	Þ



Full Screen

Close

Quit

# 2. The flag graph



## $BS(\mathcal{M})$ and $Co(\mathcal{M})$ .

A map is orientable if and only if its flag graph is bipartite.

●First ●Prev ●Next ●Last ●Go Back ●Full Screen ●Close ●Qui



Combinatorial Maps The flag graph Vertex splitting . . . Reduction to . . . Classification of . . . Joining two maps Regular maps on . . . The 3-block tree Matroids  $\Delta$ -matroids Home Page Title Page •• •• ▶ Page 7 of 36



Quit







3.

Combinatorial Maps	
The flag graph	
Vertex splitting	
Reduction to	
Classification of	
Joining two maps	
Regular maps on	
The 3-block tree	
Matroids	
$\Delta$ -matroids	
Home Page	
I itle Page	







Quit

# Vertex splitting and edge contraction



Vertex split/edge contraction.

Number of faces stays the same. Contraction of a loop is undefined.



Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps Regular maps on... The 3-block tree Matroids Δ-matroids





Close

Quit

## 4. Reduction to unitary maps

A map is called *unitary* if it has exactly one vertex and one face. By a sequence of edge contractions one can reduce the number of vertices of a map M, or the number of faces of  $M^*$ .

The flags  $\{x, \tau_0 x, \tau_2 x, \mathcal{E}x\}$  of a unitary map are distributed among the two disjoint cycles of  $\mathcal{V}$  by

$$(x, A, \mathcal{E}x, B)(\tau_2 B, \tau_0 x, \tau_2 A, \tau_2 x)$$

or

$$(x, A, \tau_0 x, B)(\tau_2 B, \mathcal{E}x, \tau_2 A, \tau_2 x)$$

Crosscap

A crosscap is assembled if A or B is empty. Crosscaps may be assembled by a sequence of vertex splits and edge contractions.



5

Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps Regular maps on... The 3-block tree Matroids  $\Delta$ -matroids





Close

Quit

		•		C		· · · · · ·	<b>P</b>
. (	Classi	ficati	on c	<b>)</b>	5	ur	taces

5.1. Normal forms



Map projection and polygon models of canonical normal forms.

**Theorem 1** Every closed surface has the topological type of either

- 1. The sphere.  $(\chi(S) = 2)$ .
- 2. A connected sum of n tori.  $(\chi(T^n) = 2(n-1)).$
- 3. A connected sum of n projective planes.  $(\chi(P^n) = 2-n)$ .

5



#### Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps

Regular maps on . . .

The 3-block tree

Matroids

 $\Delta$ -matroids

Home Page



••	••
•	

Page **11** of <mark>36</mark>



Full Screen

Close

Quit

## 6. Joining two maps

### 6.1. Connected sum



The connected sum of two tori.

$$\chi(S \sharp S') = \chi(S) + \chi(S') - 2.$$

●First ●Prev ●Next ●Last ●Go Back ●Full Screen ●Close ●Quit



#### Combinatorial Maps

- The flag graph
- Vertex splitting . . .
- Reduction to . . .
- Classification of . . .
- Joining two maps
- Regular maps on . . .
- The 3-block tree
- Matroids
- $\Delta$ -matroids
  - Home Page
  - Title Page
- ↓
  ↓
  ↓







Close

Quit

### 6.2. Edge join



 $\mathcal{M}_1 +_{x_1,y_1} \mathcal{M}_2$ , joining two maps along an edge.



#### Combinatorial Maps The flag graph

- Vertex splitting . . .
- Reduction to . . .
- Classification of . . .
- Joining two maps
- Regular maps on . . .
- The 3-block tree
- Matroids
- $\Delta$ -matroids
  - Home Page
  - Title Page
- ↓
   ↓
   Page 13 of 36
- Go Back Full Screen
  - Close
  - Quit

### 6.3. Vertex join



Two ways to join two maps at a vertex. Observe the sequence of vertex valences around the exterior face.



#### Combinatorial Maps

- The flag graph
- Vertex splitting . . .
- Reduction to . . .
- Classification of . . .
- Joining two maps
- Regular maps on . . .
- The 3-block tree
- Matroids
- $\Delta$ -matroids

Home Page

Title Page

••	••
•	Þ
Page 1	4 of 36

Full Screen

Close

Quit



A Whitney twist:  $\mathcal{M}_1 +_{x_1,\tau_2 y_1} \mathcal{M}_2$ 



- Combinatorial Maps
- The flag graph
- Vertex splitting . . .
- Reduction to . . .
- Classification of . . .
- Joining two maps
- Regular maps on . . .
- The 3-block tree
- Matroids
- $\Delta$ -matroids
  - Home Page
  - Title Page
- ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
   ✓
- Go Back
- Full Screen
  - Close
  - Quit



A Whitney flip:  $\mathcal{M}_1 +_{x_1,\tau_0 y_1} \mathcal{M}_2$ .



Combinatorial Maps The flag graph Vertex splitting ... Reduction to . . . Classification of ... Joining two maps Regular maps on . . . The 3-block tree Matroids  $\Delta$ -matroids



Page 16 of 36	
Go Back	

Close

Quit

#### **Regular** maps on the sphere 7.

If M is a regular map, then the flag graph is a Cayley graph of Aut(M) where the generating set S has three elements, namely the three group elements which map a given flag  $\phi$  to  $\tau_0 \phi$ ,  $\tau_1 \phi$ , and  $\tau_2 \phi$ .



Constructing a Cayley map for the icosahedron.



Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps Regular maps on... The 3-block tree Matroids Δ-matroids



Full Screen

Close

Quit

## 7.1. Straightening Lemma

A complex  $\mathcal{M}$  is *harmonious* if  $Aut\mathcal{M} = \text{Isom}(\mathcal{M})$ . In [6] it was shown that every planar combinatorial map can be straightened to a harmonious map.

**Theorem 2** Let  $\mathcal{M}$  be a complex of polygons piecewise linearly embedded on the sphere  $S^2$ , the Euclidean plane  $\mathbb{R}^2$ , or the hyperbolic plane  $\mathbb{H}^2$ , and  $\mathcal{M}$  is the associated map. If Aut  $\mathcal{M}$  has finitely many face orbits, then there is a complex Ideal( $\mathcal{M}$ ) of  $\mathbb{S}^2$ ,  $\mathbb{E}^2$  or  $\mathbb{H}^2$  which is combinatorially isomorphic to  $\mathcal{M}$  and such that every automorphism of  $\mathcal{M}$  is expressed as an isometry of Ideal( $\mathcal{M}$ ), so  $Isom(Ideal(\mathcal{M})) = Aut(\mathcal{M})$ .



Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of...

Joining two maps

Regular maps on . . .

The 3-block tree

Matroids

 $\Delta$ -matroids



Page **18** of **36** 

Go Back

Full Screen

Close

Quit

(pm, pm)b.

(pm, pm)a.

**Theorem 3** Every self-dual map  $\mathcal{M}$  with finitely many face orbits is the map of a harmoniously self-dual complex.





●First ●Prev ●Next ●Last ●Go Back ●Full Screen ●Close ●Qu



Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps Regular maps on... The 3-block tree Matroids



Close

Quit

## 8. The 3-block tree

For planar graphs which are 3-connected, Whitney's theorem [9] says that all its planar or spherical maps are combinatorially isomorphic and therefore, by the straightening lemma, can be studied with geometric methods alone.

For graphs of lower connectivity, faces are not uniquely defined.



The 3-block tree.



Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps Regular maps on... The 3-block tree Matroids Δ-matroids



Go Back	
Full Screen	
Close	
Quit	

For a planar graph G, we may now analyze the automorphisms of its 3-blocks via geometric methods, and then piece these symmetries together combinatorially via its structure trees. This program has been followed for planar Cayley graphs [3] and self-dual graphs and matroids, see [6] and [7], where the construction of all self-dual graphs are described, even those which have no associated self-dual map to be straightened.



A self-dual graph drawn on an unfolded cube with no corresponding self-dual map.

A refinement of the 24 symmetry pairings of maps on the sphere was obtained by Graver and Hartung in [5] by considering grid patches.



- Combinatorial Maps The flag graph Vertex splitting ... Reduction to .... Classification of ... Joining two maps Regular maps on . . . The 3-block tree
- Home Page Title Page 44 Page 21 of 36 Go Back Full Screen

Close

## Matroids

 $\Delta$ -matroids

Quit

### Matroids 9.

Whitney [10] defined a matroid on a finite set E to be a pair  $(E, \mathcal{I})$  where E is a finite set and  $\mathcal{I}$  is a collection of subsets of E such that



**I2** If  $I_1 \in \mathcal{I}$  and  $I_2 \subset I_1$ , then  $I_2 \in \mathcal{I}$ 

**I3** If  $I_1$  and  $I_2$  are members of  $\mathcal{I}$  and  $|I_1| < |I_2|$ , then there exists an element e in  $I_2 - I_1$  such that  $I_1 + e$  is a member of  $\mathcal{I}$ .



Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps Regular maps on... The 3-block tree Matroids

Home Page
Title Page
•• ••
Page 22 of 36
Go Back
Full Screen

Close

Quit

Because of condition [I2], all of the maximal independent sets have the same cardinality. These maximal independent sets are called the *bases* of the matroid. The bases may be described directly: Let E be a finite set, a nonempty collection  $\mathcal{B}$  of subsets of E is called a *basis system* for E if

B1  $\mathcal{B} \neq \emptyset$ 

**B2** For all  $B_1, B_2 \in \mathcal{B}, |B_1| = |B_2|$ 

**B3** For all  $B_1, B_2 \in \mathcal{B}$  and  $e_1 \in B_1 - B_2$ , there exists  $e_2 \in B_2 - B_1$  such that  $B_1 - e_1 + e_2 \in \mathcal{B}$ .

Condition [B3] is sometimes called the *exchange axiom*. It also has a slightly different but equivalent formulation:

**B3'** For all  $B_1, B_2 \in \mathcal{B}$  and  $e_2 \in B_2 - B_1$ , there exists  $e_1 \in B_1 - B_2$  such that  $B_1 - e_1 + e_2 \in \mathcal{B}$ .

Complements of bases also satisfy [B3], these complements are bases of the dual matroid. Every matroid M has a dual  $M^*$ .



```
Combinatorial Maps
The flag graph
Vertex splitting . . . .
Reduction to . . . .
Classification of . . . .
Joining two maps
Regular maps on . . .
The 3-block tree
Matroids
A-matroids
```

```
Home Page
Title Page
```

```
Full Screen
```

Close

Quit

Let M be a matroid on E with independent sets  $\mathcal{I}$  and define  $r_{(\mathcal{I})}$ , a function from the power set of E into the nonnegative integers by  $r_{(\mathcal{I})}(S) = \max\{|I| : I \in \mathcal{I}, I \subseteq S\}$ . The function  $r = r_{\mathcal{I}}$  is called the *rank function of* M.

In general, let E be a finite set and r a function from the power set of E into the nonnegative integers so that

**R1** 
$$r(\emptyset) = 0;$$

```
\mathbf{R2} \ r(S) \le |S|;
```

**R3** if  $S \subseteq T$  then  $r(S) \leq r(T)$ ;

 $\mathbf{R4}\ r(S\cup T)+r(S\cap T)\leq r(S)+r(T);$ 

then r is called a rank function on E. If r is a rank function on E we define  $\mathcal{I}(r) = \{I \subseteq E \mid r(I) = |I|\}$ Condition [R4] is called the submodular inequality.



- Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps Regular maps on... The 3-block tree Matroids

Close

 $\Delta$ -matroids

CI

Quit

Given a finite set E, we call a collection C a *cycle system* for E, if the following three conditions are satisfied:

**Z1** If  $C \in \mathcal{C}$  then  $C \neq \emptyset$ 

**Z2** If  $C_1$  and  $C_2$  are members of  $\mathcal{C}$  then  $C_1 \not\subseteq C_2$ 

**Z3** If  $C_1$  and  $C_2$  are members of  $\mathcal{C}$  and if e' is an element of  $C_1 \cap C_2$ , then for each  $e \in (C_2 - C_1)$  there is an element  $C \in \mathcal{C}$ , such that  $e \in C \subseteq (C_1 \cap C_2 - e')$ .



Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of ... Joining two maps Regular maps on... The 3-block tree Matroids Δ-matroids



Full Screen

Close

Quit

A matroid is graphic if it is isomorphic to the cycle matroid on a graph G. Non-isomorphic graphs may have the same cycle matroid, but 3-connected graphs are uniquely determined by their matroids.

M is co-graphic if  $M^*$  is graphic.

M is graphic as well as co-graphic iff G is planar. Map duality (geometric duality) agrees with matroid duality. If G(V, E) is planar and connected, its cycle matroid has rank |V| - 1, its co-cycle matroid has rank |F| - 1, so |V| - 1 + |F| - 1 = |E|. The facial cycles generate the cycle space.



Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps Regular maps on... The 3-block tree Matroids Δ-matroids

Home Page
Title Page

• •

Page 26 of 36

Go Back

Full Screen

Close

Quit



cubeflag01.pdf



Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps Regular maps on... The 3-block tree Matroids



Full Screen

Close

Quit

## **10.** $\triangle$ -matroids

Introduced by Bouchet as set systems satisfying the symmetric exchange axiom

For  $F', F'' \in \mathcal{F}, x \in F' \Delta F''$ , there exists  $y \in F'' \Delta F'$  such that  $F' \Delta \{x, y\} \in \mathcal{F}$ .

Feasible sets need not be equicardinal.

In [2] Bouchet associates a  $\Delta$ -matroid to a map on a topological surface S by defining edge sets F feasible if  $S - cl(F \cup \overline{F}^*)$  is connected. This easily translates to connectivity properties of the flag graph.



Combinatorial Maps The flag graph Vertex splitting . . . . Reduction to . . . . Classification of . . . . Joining two maps Regular maps on . . . . Regular maps on . . . . The 3-block tree Matroids A-matroids Home Page



Page 28 of 36

Go Back

Full Screen

Close

Quit

### For example consider $K_4$ embedded on a torus.



●First ●Prev ●Next ●Last ●Go Back ●Full Screen ●Close ●Quit



- Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps Regular maps on... The 3-block tree
- Matroids
- $\Delta$ -matroids



Close

Quit





Combinatorial Maps The flag graph Vertex splitting . . . Reduction to .... Classification of ... Joining two maps Regular maps on . . . The 3-block tree Matroids  $\Delta$ -matroids Home Page Title Page •• 44 Page **30** of **36** Go Back Full Screen

Close

Quit

From each red-green square both edges of one color must be deleted without destroying connectivity of the flag graph.





- Combinatorial Maps The flag graph Vertex splitting... Reduction to... Classification of... Joining two maps Regular maps on... The 3-block tree
- Matroids
- $\Delta$ -matroids

Home Page	
Title Page	
•• ••	
<ul> <li>↓</li> </ul>	
Page 31 of 36	
Go Back	
Full Screen	

Close

Quit

**Theorem 4** Let D(M) be the  $\Delta$ -matroid of a map M(G, S)and let  $M^*(G^*, S)$  be the dual map. Then

- $\bullet \ D(M^*) = D(M)^*;$
- the lower matroid of D is the cycle matroid of G;
- the upper matroid of D is the co-cycle matroid of  $G^*$ ;

• 
$$w(D) = 2 - \chi(S).$$





Note that there are connections to 2-matroids and ribbon graphs. A good reference for ribbon graphs is [4], see also [1]. f





Close

Quit

A ribbon graph.



#### Combinatorial Maps The flag graph Vertex splitting . . . . Reduction to . . . Classification of . . . . Joining two maps Regular maps on . . . The 3-block tree Matroids A-matroids





Close

Quit

 $K_5$  on the torus





●First ●Prev ●Next ●Last ●Go Back ●Full Screen ●Close ●Qui



- Combinatorial Maps The flag graph Vertex splitting . . . Reduction to . . . Classification of . . . Joining two maps Regular maps on . . .
- The 3-block tree
- Matroids
- $\Delta$ -matroids



Full Screen

Close

Quit

### Another way to embed $K_5$ on the torus



●First ●Prev ●Next ●Last ●Go Back ●Full Screen ●Close ●Qu



- Combinatorial Maps The flag graph Vertex splitting...
- Reduction to . . .
- Classification of . . .
- Joining two maps
- Regular maps on . . .
- The 3-block tree
- Matroids
- $\Delta$ -matroids





Full Screen

Close









### References

- [1] Remi Cocou Avohou. Polynomial invariants for arbitrary rank D weakly-colored stranded graphs. SIGMA Symmetry Integrability Geom. Methods Appl., 12:Paper No. 030, 23, 2016.
- [2] André Bouchet. Maps and  $\triangle$ -matroids. Discrete Math., 78(1-2):59–71, 1989.
- [3] Carl Droms, Brigitte Servatius, and Herman Servatius. Connectivity and planarity of Cayley graphs. *Beiträge Algebra Geom.*, 39(2):269–282, 1998.
- [4] Joanna A. Ellis-Monaghan and Iain Moffatt. *Graphs on surfaces*. SpringerBriefs in Mathematics. Springer, New York, 2013. Dualities, polynomials, and knots.
- [5] Jack E. Graver and Elizabeth J. Hartung. Self-dual spherical grids. *Electron. J. Combin.*, 21(1):Paper 1.36, 36, 2014.
- Brigitte Servatius and Herman Servatius. The 24 symmetry pairings of self-dual maps on the sphere. Discrete Math., 140(1-3):167–183, 1995.
- [7] Brigitte Servatius and Herman Servatius. Self-dual graphs. Discrete Math., 149(1-3):223-232, 1996.
- [8] William T. Tutte. What is a map? In New directions in the theory of graphs (Proc. Third Ann Arbor Conf., Univ. Michigan, Ann Arbor, Mich., 1971), pages 309–325. Academic Press, New York, 1973.
- [9] Hassler Whitney. Congruent Graphs and the Connectivity of Graphs. Amer. J. Math., 54(1):150–168, 1932.
- [10] Hassler Whitney. On the Abstract Properties of Linear Dependence. Amer. J. Math., 57(3):509– 533, 1935.



Combinatorial Maps

- The flag graph
- Vertex splitting . . .
- Reduction to . . .
- Classification of . .
- Joining two maps
- Regular maps on . . .
- The 3-block tree
- Matroids
- $\Delta$ -matroids





Go Back



Close

