



Combinatorial Rigidity and the Molecular Conjecture

Brigitte Servatius
Worcester Polytechnic Institute


The Proof of the Product Rule


To derivate a product as defined
The diff'rence quotient limit do compute,
And in that simple limit try to find
Each factor's diff'rence quotient, how astute!

But products of the functions at each x
Do not combine or factor, not on cue –
And fudging it does not survive the checks! :-(
A bit of clever insight finds the clue,

A product term with x 's, ah, distinct
With plus and minus versions introduced.
The limit now with algebra succinct
Computes, so thus that rule can be deduced: :-)


Add factor one times factor two derived
To factor two times factor one derived.



[Introduction](#)[Type of Rigidity](#)[Vertex transitive...](#)[Example](#)[Random graphs](#)[More general...](#)[Modeling molecules](#)[Molecular...](#)[Realization in the...](#)[Combinatorial...](#)[Open problems](#)[Home Page](#)[Title Page](#)[Page 2 of 65](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Problem

Suppose we want to build a rigid structure



such as a **GREENHOUSE**.

Do we hire an architect and design it? [Answer: **YES**.]

How does the architect determine if the structure is rigid?

[Introduction](#)[Type of Rigidity](#)[Vertex transitive...](#)[Example](#)[Random graphs](#)[More general...](#)[Modeling molecules](#)[Molecular...](#)[Realization in the...](#)[Combinatorial...](#)[Open problems](#)[Home Page](#)[Title Page](#)[Page 3 of 65](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

1. Introduction

Generic rigidity in the plane is a graph theoretic property:

Theorem 1 (*Laman - 1970*)[7] *A graph is generically rigid in the plane if and only if it has a subset $|E|$ of edges with*

$$|E| = 2|V(E)| - 3$$

and, for every subset $F \subseteq E$,

$$|F| \leq 2|V(F)| - 3.$$



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀ ▶

◀ ▶


Page 4 of 65

Go Back

Full Screen

Close

Quit



Rigidity
Different notions of rigidity.



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀ ▶

◀ ▶


Page 5 of 65

Go Back

Full Screen

Close

Quit



Rigidity
Non rigid graphs have a motion.



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀ ▶

◀ ▶


Page 6 of 65

Go Back

Full Screen

Close

Quit



Non-infinitesimally rigid graphs have initial velocity candidates.



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀ ▶

◀ ▶


Page 7 of 65

Go Back

Full Screen

Close

Quit



Generic rigidity is a property of the graph,
not the embedding.



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀◀ ▶▶

◀ ▶

Page 8 of 65

Go Back

Full Screen

Close

Quit

2. Type of Rigidity

We will use the term *framework* (in m -space) to denote a triple (V, E, \vec{p}) , where (V, E) is a graph and \vec{p} is an embedding (injection) of V into real m -space.



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀ ▶

◀ ▶

Page 9 of 65

Go Back


Full Screen

Close

Quit

We say that a framework is *globally rigid* (in m -space) if all solutions to the system of quadratic equations obtained from requiring all edge lengths to be fixed, with the coordinates of the vertices as variables, correspond to congruent frameworks; we say that a framework is *rigid* (in m -space) if all solutions to the corresponding system in some neighborhood of the original solution (as a point in mn -space) come from congruent frameworks.

DON'T CLICK HERE!



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀◀ ▶▶

◀ ▶

Page 10 of 65

Go Back

Full Screen

Close

Quit

We say that a given framework $(V, E, \overrightarrow{p})$ is *generic* if all frameworks corresponding to points in a neighborhood of $P = \overrightarrow{p}(V)$ in \mathbb{R}^{nm} are rigid or not rigid as is $(V, E, \overrightarrow{p})$. A set of points P in m -space is said to be *generic* if each framework $(V, E, \overrightarrow{p})$ with $\overrightarrow{p}(V) = P$ is generic.



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀ ▶

◀ ▶

Page 11 of 65

Go Back

Full Screen

Close

Quit

If $G(V, E)$ is not rigid, we call the maximal rigid subgraphs of G the *rigid components* and note that rigid components partition E . Then $\mathcal{M}(G)$ is the direct sum over its restrictions on the rigid components.

The following theorem is equivalent to Laman's Theorem 1, it uses the rank function of \mathcal{M} rather than independence to characterize rigidity.



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀◀ ▶▶

◀ ▶

Page 12 of 65

Go Back

Full Screen

Close

Quit

Theorem 2 [9] *Let $G = (V, E)$ be a graph. Then G is rigid if and only if for all families of induced subgraphs $\{G_i = (V_i, E_i)\}_{i=1}^m$ such that $E = \cup_{i=1}^m E_i$ we have $\sum_{i=1}^m (2|V_i| - 3) \geq 2|V| - 3$.*



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀ ▶

◀ ▶

Page 13 of 65

Go Back

Full Screen

Close

Quit

$G(V, E)$ is called *redundantly rigid* if $G(V, E - e)$ is rigid for all $e \in E$, i.e. the removal of a single edge e from the rigid graph G does not destroy rigidity. Redundant rigidity is a key to characterize global rigidity.

Theorem 3 [5] *Let G be a graph. Then G is globally rigid if and only if G is a complete graph on at most three vertices, or G is both 3-connected and redundantly rigid.*



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀▶

◀▶

Page 14 of 65

Go Back

Full Screen

Close

Quit

3. Vertex transitive graphs

Theorem 4 *A four-regular vertex transitive graph is generically rigid in the plane if and only if it contains no subgraph isomorphic to K_4 , or is K_5 or one of the graphs in the following figure.*



Introduction

Type of Rigidity

Vertex transitive...

Example

Random graphs

More general...

Modeling molecules

Molecular...

Realization in the...

Combinatorial...

Open problems

Home Page

Title Page



Page 15 of 65

Go Back

Full Screen


Close

Quit



$K_4 \times K_2$

a)



b)

Vertex transitive rigid graphs containing K_4 .



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀◀

▶▶

◀

▶

Page 16 of 65

Go Back

Full Screen

Close

Quit

Theorem 5 *Let G be a connected k -regular vertex transitive graph on n vertices. Then G is not rigid if and only if either:*

(a) $k = 2$ and $n \geq 4$.

(b) $k = 3$ and $n \geq 8$.

(c) $k = 4$ and G has a factor consisting of s disjoint copies of K_4 where $s \geq 4$

(d) $k = 5$ and G has a factor consisting of t disjoint copies of K_5 where $t \geq 8$.



Introduction

Type of Rigidity

Vertex transitive...

Example

Random graphs

More general...

Modeling molecules

Molecular...

Realization in the...

Combinatorial...

Open problems

Home Page

Title Page





Page 17 of 65

Go Back



Full Screen

Close


Quit



Two embeddings which are rigid, but neither infinitesimally rigid nor globally rigid.



Two embeddings which are rigid and infinitesimally rigid but not globally rigid.



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀ ▶

◀ ▶


Page 18 of 65

Go Back


Full Screen

Close


Quit



Two embeddings which are rigid, but not globally rigid.



4. Example



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

[Home Page](#)

[Title Page](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 19 of 65

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)



[Introduction](#)

[Type of Rigidity](#)

[Vertex transitive...](#)

[Example](#)

[Random graphs](#)

[More general...](#)

[Modeling molecules](#)

[Molecular...](#)

[Realization in the...](#)

[Combinatorial...](#)

[Open problems](#)

[Home Page](#)

[Title Page](#)




Page 20 of 65

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)





[Introduction](#)

[Type of Rigidity](#)

[Vertex transitive...](#)

[Example](#)

[Random graphs](#)

[More general...](#)

[Modeling molecules](#)

[Molecular...](#)

[Realization in the...](#)

[Combinatorial...](#)

[Open problems](#)

[Home Page](#)

[Title Page](#)




Page 21 of 65

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)





[Introduction](#)

[Type of Rigidity](#)

[Vertex transitive...](#)

[Example](#)

[Random graphs](#)

[More general...](#)

[Modeling molecules](#)

[Molecular...](#)

[Realization in the...](#)

[Combinatorial...](#)

[Open problems](#)

[Home Page](#)

[Title Page](#)




Page 22 of 65

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)





[Introduction](#)

[Type of Rigidity](#)

[Vertex transitive...](#)

[Example](#)

[Random graphs](#)

[More general...](#)

[Modeling molecules](#)

[Molecular...](#)

[Realization in the...](#)

[Combinatorial...](#)

[Open problems](#)

[Home Page](#)

[Title Page](#)




Page 23 of 65

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)





[Introduction](#)

[Type of Rigidity](#)

[Vertex transitive...](#)

[Example](#)

[Random graphs](#)

[More general...](#)

[Modeling molecules](#)

[Molecular...](#)

[Realization in the...](#)

[Combinatorial...](#)

[Open problems](#)

[Home Page](#)

[Title Page](#)




Page 24 of 65

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)





[Introduction](#)

[Type of Rigidity](#)

[Vertex transitive...](#)

[Example](#)

[Random graphs](#)

[More general...](#)

[Modeling molecules](#)

[Molecular...](#)

[Realization in the...](#)

[Combinatorial...](#)

[Open problems](#)

[Home Page](#)

[Title Page](#)




Page 25 of 65

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)





[Introduction](#)

[Type of Rigidity](#)

[Vertex transitive...](#)

[Example](#)

[Random graphs](#)

[More general...](#)

[Modeling molecules](#)

[Molecular...](#)

[Realization in the...](#)

[Combinatorial...](#)

[Open problems](#)

[Home Page](#)

[Title Page](#)




Page 26 of 65

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)





[Introduction](#)

[Type of Rigidity](#)

[Vertex transitive...](#)

[Example](#)

[Random graphs](#)

[More general...](#)

[Modeling molecules](#)

[Molecular...](#)

[Realization in the...](#)

[Combinatorial...](#)

[Open problems](#)

[Home Page](#)

[Title Page](#)




Page 27 of 65

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)





[Introduction](#)

[Type of Rigidity](#)

[Vertex transitive...](#)

[Example](#)

[Random graphs](#)

[More general...](#)

[Modeling molecules](#)

[Molecular...](#)

[Realization in the...](#)

[Combinatorial...](#)

[Open problems](#)

[Home Page](#)

[Title Page](#)




Page 28 of 65

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)





[Introduction](#)

[Type of Rigidity](#)

[Vertex transitive...](#)

[Example](#)

[Random graphs](#)

[More general...](#)

[Modeling molecules](#)

[Molecular...](#)

[Realization in the...](#)

[Combinatorial...](#)

[Open problems](#)

[Home Page](#)

[Title Page](#)




Page 29 of 65

[Go Back](#)

[Full Screen](#)


[Close](#)

[Quit](#)





Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

[Home Page](#)[Title Page](#)[Page 30 of 65](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀◀

▶▶

◀

▶

Page 31 of 65

Go Back

Full Screen

Close

Quit

However, Lovász and Yemini [9] note that their proof technique will show that $G - \{e_1, e_2, e_3\}$ is rigid for all $e_1, e_2, e_3 \in E$, and hence that G is redundantly rigid. This result was combined with Theorem 3 in [5] to deduce

Theorem 6 *Every 6-connected graph is globally rigid.*



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀ ▶

◀ ▶

Page 32 of 65

Go Back

Full Screen

Close

Quit

5. Random graphs

Let $G_{n,d}$ denote the probability space of all d -regular graphs on n vertices with the uniform probability distribution. A sequence of graph properties A_n holds asymptotically almost surely, or a.a.s. for short, in $G_{n,d}$ if $\lim_{n \rightarrow \infty} \Pr_{G_{n,d}}(A_n) = 1$. Graphs in $G_{n,d}$ are known to be a.a.s. highly connected. It was shown by Bollobás [1] and Wormald [12] that if $G \in G_{n,d}$ for any fixed $d \geq 3$, then G is a.a.s. d -connected. This result was extended to all $3 \leq d \leq n - 4$ by Cooper et al. [3] and Krivelevich et al. [6]. Stronger results hold if we discount ‘trivial’ cutsets. In [13], Wormald shows that if $G \in G_{n,d}$ for any fixed $d \geq 3$, then G is a.a.s. cyclically $(3d - 6)$ -edge-connected.



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀◀

▶▶

◀

▶

Page 33 of 65

Go Back

Full Screen

Close

Quit

Theorem 7 *If $G \in G_{n,4}$ then G is a.a.s. globally rigid.*

In fact this result holds for all $d \geq 4$.

Theorem 8 *If $G \in G_{n,d}$ and $d \geq 4$ then G is a.a.s. globally rigid.*



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀ ▶

◀ ▶

Page 34 of 65

Go Back

Full Screen

Close

Quit

Let $G(n, p)$ denote the probability space of all graphs on n vertices in which each edge is chosen independently with probability p . In the following we will assume that all logarithms are natural. We will need the following results on $G(n, p)$.

Lemma 1 *Let $G \in G(n, p)$, where $p = (\log n + k \log \log n + w(n))/n$, $k \geq 2$ is an integer and $\lim_{n \rightarrow \infty} w(n) = \infty$. For each fixed integer t , let S_t be the set of vertices of G of degree at most t . Then, a.a.s.*

- (a) S_{k-1} is empty,
- (b) no two vertices of S_t are joined by a path of length at most two in G ,
- (c) $G - S_{t-1}$ is non-empty and t -connected.

PROOF: Facts (a) and (b) are well known, see for example [2]. Fact (c) follows from (a), (b) and [10, Theorem 4] \square



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀ ▶

◀ ▶

Page 35 of 65

Go Back

Full Screen

Close

Quit

Theorem 9 Let $G \in G(n, p)$, where $p = (\log n + k \log \log n + w(n))/n$, and $\lim_{n \rightarrow \infty} w(n) = \infty$.

(a) If $k = 2$ then G is a.a.s. rigid.

(b) If $k = 3$ then G is a.a.s. globally rigid.

PROOF: (a) We adopt the notation of Lemma 1. It follows from Lemma 1 that a.a.s. $S_1 = \emptyset$ and $G - S_5$ is a.a.s. 6-connected. Hence $G - S_5$ is a.a.s. (globally) rigid by Theorem 6. Since adding a new vertex joined by at least two new edges to a rigid graph preserves rigidity, it follows that G is a.a.s. rigid.

(b) This follows in similar way to (a), using the facts that $S_2 = \emptyset$ and that adding a new vertex joined by at least three new edges to a globally rigid graph preserves global rigidity. \square



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀◀ ▶▶

◀ ▶

Page 36 of 65

Go Back

Full Screen

Close


Quit

The bounds on p given in Theorem 9 are best possible since if $G \in G(n, p)$ and $p = (\log n + k \log \log n + c)/n$ for any constant c , then G does not a.a.s. have minimum degree at least k , see [2].



The Kagome Lattice

Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems



[Home Page](#)

[Title Page](#)



Page 37 of 65

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀ ▶

◀ ▶

Page 38 of 65

Go Back

Full Screen

Close

Quit

Let $Geom(n, r)$ denote the probability space of all graphs on n vertices in which the vertices are distributed uniformly at random in the unit square and each pair of vertices of distance at most r are joined by an edge. Suppose $G \in Geom(n, r)$. Li, Wan and Wang [8] have shown that if $n\pi r^2 = \log n + (2k - 3) \log \log n + w(n)$ for $k \geq 2$ a fixed integer and $\lim_{n \rightarrow \infty} w(n) = \infty$, then G is a.a.s. k -connected. As noted by Eren et al. [4], this result can be combined with Theorem 6 to deduce that if $n\pi r^2 = \log n + 9 \log \log n + w(n)$ then G is a.a.s. globally rigid. On the other hand, it is also shown in [8] that if $n\pi r^2 = \log n + (k - 1) \log \log n + c$ for any constant c , then G is not a.a.s. k -connected. It is still conceivable, however, that if $n\pi r^2 = \log n + \log \log n + w(n)$ then G is a.a.s. rigid, and that if $n\pi r^2 = \log n + 2 \log \log n + w(n)$ then G is a.a.s. globally rigid.



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules**
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀◀

▶▶

◀

▶

Page 39 of 65

Go Back

Full Screen

Close

Quit

6. More general structures

Bar and joint frameworks

rigid bars (edges), universal joints (vertices)

Body and hinge frameworks

rigid bodies (vertices), hinges (edges)



n-space: Generic Body and Hinge Frameworks


Solved

Body and Hinge Frameworks in 3-space:

Each rigid body has 6 degrees of freedom.

If two bodies are joined along a linear hinge the resulting structure has one internal degree of freedom.

Each hinge removes 5 degrees of freedom.



Home Page

Title Page

◀◀

▶▶

◀

▶

Page 40 of 65

Go Back

Full Screen

Close

Quit



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀ ▶

◀ ▶

Page 41 of 65

Go Back

Full Screen


Close

Quit

Graph $G = (B, H)$
B: vertices for abstract bodies,
H: for pairs of bodies sharing a hinge.
Necessary condition for independence:

$$5|H'| \leq 6|B'| - 6$$

Theorem 10 (Tay and Whiteley – 1984) *The necessary condition is also sufficient for generic independence.*



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀ ▶

◀ ▶

Page 42 of 65

Go Back

Full Screen

Close

Quit

Algorithms:

$$6|B'| - 6 = 6(|B'| - 1)$$


or
6 spanning trees in $5G(B, H)$, which is the multi-graph obtained from $G(B, H)$, by replacing each edge by a set of 5 parallel edges.



7. Modeling molecules

(special graphs) - can we predict rigidity?

Single atom and associated bonds



$$|V| = 5 \quad |E| = 10$$

$$|E| = 3|V| - 5 \quad \text{overbraced}$$

Introduction

Type of Rigidity

Vertex transitive...

Example

Random graphs

More general...

Modeling molecules

Molecular...

Realization in the...

Combinatorial...

Open problems

Home Page

Title Page

◀◀

▶▶

◀

▶

Page 43 of 65

Go Back

Full Screen

Close

Quit



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀ ▶

◀ ▶

Page 44 of 65

Go Back


Full Screen

Close

Quit

Adjacent atom clusters

Flexible




$$\begin{aligned} |B| &= 2, |H| = 1, \\ |V| &= 4, |E| = 5, \end{aligned}$$

$$\begin{aligned} 5|H| &= 6|B| - 7, \\ |E| &= 3|V| - 7 \end{aligned}$$



Rings of atoms:

Ring of 6 atoms and bonds



Bar and Joint:

$$|V| = 6, |E| = 12, |E| = 3|V| - 6$$

Body and hinge:

$$|B| = 6, |H| = 6, 5|H| = 6|B| - 6$$

Just the right number to be rigid - generically.

- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀ ▶

◀ ▶

Page 45 of 65

Go Back

Full Screen

Close

Quit



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

[Home Page](#)[Title Page](#)

Page 46 of 65

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Graph G of atoms and covalent bonds

Can we use the body and hinge model to predict rigidity?

Problem: Some hinges are concurrent

Special geometry may lower rank!



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page



Page 47 of 65

Go Back

Full Screen

Close

Quit

Graph G of atoms and covalent bonds

To utilize the bar and joint model, form G^2 :

The new edges model second neighbor bond bending pairs.

Count as

$$3|V| - 6$$

priority system on bond edges.

Problem: for general graphs G the rank may be lower. (May work for G^2 ?)



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

[Home Page](#)

[Title Page](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 48 of 65

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

Lots of experimental evidence

Proofs of correctness for special classes of graphs

Plausibility arguments related to other conjectures on 3-space rigidity

Sketched proof of equivalence of the two conjectures.

Conjectures embedded in implemented algorithms: FIRST on the web (Arizona State University)



8. Molecular Conjecture in the Plane

Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page



Page 49 of 65

Go Back

Full Screen

Close

Quit


Given: Simple graph $G = (V, E)$.

- Regard G as a body and pin graph of a structure in the plane:

Vertices are bodies.

Edges denote pins.


- Note: Each pin connects just two bodies. Otherwise we would need a hyper-graph.
- Realizations:
 - Amorphous bodies. Embedding specifies the location of the pins.
 - Line bodies. Embedding may specify either lines or pins.
 - Question: Does the line realization always exist?



9. Realization in the Plane.

Theorem 11 *If $G = (V, E)$ is simple, then a pin collinear structure exists.*

Take any generic embedding of the structure graph $G = (V, E)$ in \mathbb{R}^2 . Form the polar of that embedding.

[Home Page](#)[Title Page](#)[◀](#)[▶](#)[◀](#)[▶](#)[Page 50 of 65](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)




Question:

Is the polar generic as a line-pin structure?


Question:

Does it have the same rank as a generic body-pin structure?





- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems



Home Page

Title Page

◀ ▶

◀ ▶

Page 52 of 65

Go Back


Full Screen

Close

Quit



A general body-pin structure:



The incidence structure is a hyper-graph. Does it have a pin collinear realization?



[Introduction](#)

[Type of Rigidity](#)

[Vertex transitive...](#)

[Example](#)

[Random graphs](#)

[More general...](#)

[Modeling molecules](#)

[Molecular...](#)

[Realization in the...](#)

[Combinatorial...](#)

[Open problems](#)

[Home Page](#)

[Title Page](#)




Page 54 of 65

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)





Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀◀ ▶▶

◀ ▶

Page 55 of 65

Go Back

Full Screen

Close

Quit

Theorem 12 *A multigraph G can be realized as an infinitesimally rigid body and hinge framework in \mathbb{R}^d if and only if $((\binom{d+1}{2} - 1)G$ has $\binom{d+1}{2}$ edge-disjoint spanning trees. (Tay and Whiteley, 1984)*

[Introduction](#)[Type of Rigidity](#)[Vertex transitive...](#)[Example](#)[Random graphs](#)[More general...](#)[Modeling molecules](#)[Molecular...](#)[Realization in the...](#)[Combinatorial...](#)[Open problems](#)[Home Page](#)[Title Page](#)[◀](#)[▶](#)[◀](#)[▶](#)[Page 56 of 65](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Recent Advances in the Generic Rigidity of Structures, Tiong-Seng Tay and Walter Whiteley Structural Topology # 9, 1984
Many body and hinge structures are built under additional constraints. For example in architecture flat panels may be used in which all hinges are coplanar. In molecular chemistry, we can model molecules by rigid atoms hinged along the bond lines so that all hinges to an atom are concurrent. This is the natural projective dual for the architectural condition.

Conjecture: A multigraph is generically rigid for hinged structures in n -space iff it is generically rigid for hinged structures in n -space with all hinges of body v_i in a hyperplane H_i of the space.



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page



Page 57 of 65

Go Back

Full Screen

Close

Quit

Jackson and Jordan show that the body-and-pin and rod-and-pin 2-polymatroids of a graph are identical. As a solution to the molecular conjecture they formulate

Theorem 13 *Let $G(V, E)$ be a multigraph. Then the following statements are equivalent:*

(a) *G has a realization as an infinitesimally rigid body and hinge framework in \mathbb{R}^2 .*

(b) *G has a realization as an infinitesimally rigid body-and-hinge framework (G, q) in \mathbb{R}^2 with each of the sets of points $\{q(e) : e \in E_G(v)\}, v \in V$, collinear.*

(c) *$2G$ contains 3 edge disjoint spanning trees.*



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page

◀ ▶

◀ ▶


Page 58 of 65

Go Back

Full Screen

Close

Quit





- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀▶

◀▶

Page 59 of 65

Go Back

Full Screen

Close

Quit

$$\begin{bmatrix}
 & V & X_3 & X_2 & X_1 & Y_1 & Y_2 & Y_3 & W \\
 & & & & r_1 & -r_1 & & & \\
 & & & r_2 & & & -r_2 & & \\
 & & r_3 & & & & & -r_3 & \\
 & -x & x & & & & & & \\
 & & -x & x & & & & & \\
 & & & & -y & y & & & \\
 & & & & & -y & y & & \\
 -p_1 & & & & p_1 & & & & \\
 -p_2 & & & p_2 & & & & & \\
 -p_3 & p_3 & & & & & & & \\
 & & & & & q_1 & & & -q_1 \\
 & & & & & & q_2 & & -q_2 \\
 & & & & & & & q_3 & -q_3
 \end{bmatrix}$$



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀ ▶

◀ ▶

Page 60 of 65

Go Back

Full Screen

Close

Quit

$$\begin{array}{cccccccc}
 V & X_3 & X_2 & X_1 & Y_1 & Y_2 & Y_3 & W \\
 & & & r_1^\perp & -r_1^\perp & & & \\
 & & & r_1 & -r_1 & & & \\
 & & r_2 & & & -r_2 & & \\
 & r_3 & & & & & & -r_3 \\
 -x & x & & & & & & \\
 & -x & x & & & & & \\
 & & & & -y & y & & \\
 & & & & & -y & y & \\
 -p_1 & & & p_1 & & & & \\
 -p_2 & & p_2 & & & & & \\
 -p_3 & p_3 & & & & & & \\
 & & & & q_1 & & & -q_1 \\
 & & & & & q_2 & & -q_2 \\
 & & & & & & q_3 & -q_3
 \end{array}$$



- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems

Home Page

Title Page

◀ ▶

◀ ▶

Page 61 of 65

Go Back

Full Screen

Close




Quit

$$\begin{bmatrix}
 V & X_3 & X_2 & Z' & Z & Y_2 & Y_3 & W \\
 & & & r_1^\perp & & & & \\
 & & & r_1 & & & & \\
 & & r_2 & & & -r_2 & & \\
 & r_3 & & & & & -r_3 & \\
 -x & x & & & & & & \\
 & & -x & & x & & & \\
 & & & & -y & y & & \\
 & & & & & -y & y & \\
 -p_1 & & & & p_1 & & & \\
 -p_2 & & p_2 & & & & & \\
 -p_3 & p_3 & & & & & & \\
 & & & & q_1 & & & -q_1 \\
 & & & & & q_2 & & -q_2 \\
 & & & & & & q_3 & -q_3
 \end{bmatrix}$$



10. Combinatorial Allostery

- Introduction
- Type of Rigidity
- Vertex transitive...
- Example
- Random graphs
- More general...
- Modeling molecules
- Molecular...
- Realization in the...
- Combinatorial...
- Open problems



Home Page

Title Page

◀ ▶

◀ ▶

Page 62 of 65

Go Back

Full Screen

Close

Quit



Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

[Home Page](#)[Title Page](#)[Page 63 of 65](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

11. Open problems

- Is the solution of the molecular conjecture useful for computational biology?
- Translate combinatorial allostery to molecules.
- Are random 6-regular graphs rigid in 3-space?
- Generalizations to tensegrities?
- Can sparse random graphs be realized as unit distance graphs?



References

- [1] B. BOLLOBÁS, Random graphs, in *Surveys in Combinatorics 1999* (N.V. Temperley, ed.), LMS Lecture Note Series 52, Cambridge University Press, Cambridge, UK, 1981, 80–102.
- [2] B. BOLLOBÁS, *Random graphs* Academic Press, New York, 1985.
- [3] C. COOPER, A. FRIEZE AND B. REED, Random regular graphs of non-constant degree, *SIAM J. Algebraic Discrete Methods* 3 (1982), no. 1, 91–98.
- [4] T. EREN, D.K. GOLDENBERG, W. WHITELEY AND Y.R. YANG, Rigidity, computation, and randomization in network localization, preprint.
- [5] B. JACKSON AND T. JORDÁN, Connected rigidity matroids and unique realizations of graphs, *J. Combin. Theory Ser. B* 94 (2005), no. 1, 1–29.
- [6] M. KRIVELEVITCH, B. SUDAKOV, V.H. VU AND N.C. WORMALD, Random regular graphs of high degree, *SIAM J. Algebraic Discrete Methods* 3 (1982), no. 1, 91–98.
- [7] G.LAMAN On Graphs and rigidity of plane skeletal structures, *J. Engrg. Math.* 4 (1970), 331–340.
- [8] X-Y. LI, P-J. WAN AND Y. WANG, Fault tolerance deployment and topology control in wireless networks, in *Proceedings of the ACM Symposium on Mobile Ad Hoc Networking and Computing (Mobihoc)* (2003), Annapolis, MD, June 2003.
- [9] L. LOVÁSZ AND Y. YEMINI, On generic rigidity in the plane, *SIAM J. Algebraic Discrete Methods* 3 (1982), no. 1, 91–98.
- [10] T. LUCZAK, Size and connectivity of the k -core of a random graph, *Discrete Math.* 91 (1991), 61–68.
- [11] J.GRAVER, B. SERVATIUS AND H.SERVATIUS *Combinatorial rigidity*, American Mathematical Society (1993).



- [12] N.C. WORMALD, The asymptotic connectivity of labelled regular graphs, *J. Combinatorial Th., Series B*, (1981), 156–167.
- [13] N.C. WORMALD, Models of random regular graphs, in *Surveys in Combinatorics 1999* (J.D. Lamb and D.A. Preece, eds.), (1999), 239–298.

Introduction
Type of Rigidity
Vertex transitive...
Example
Random graphs
More general...
Modeling molecules
Molecular...
Realization in the...
Combinatorial...
Open problems

Home Page

Title Page



Page 65 of 65

Go Back

Full Screen

Close

Quit