

Confidence Interval Guide

I. Confidence Interval for a Population Mean

Assumptions

1. There are n observations from a $N(\mu, \sigma^2)$ population (normality is not crucial if sample size is large enough: 30 or more).
2. σ^2 is unknown.

Formula

A level L confidence interval for μ is $\bar{y} \pm t_{n-1, (1+L)/2} \frac{s}{\sqrt{n}}$, where \bar{y} is the sample mean and s is the sample standard deviation.

II. An Approximate Score Confidence Interval for a Population Proportion

Assumptions

1. We are interested in estimating the proportion p having a certain characteristic in the target population.
2. y is the number having the characteristic in a random sample of size n taken from the population.

Formula

An approximate score level L confidence interval for p is $\tilde{p} \pm z_{(1+L)/2} \sqrt{\tilde{p}(1-\tilde{p})/n}$, where

$$\tilde{p} = \frac{y + 0.5z_{(1+L)/2}^2}{n + z_{(1+L)/2}^2}.$$

III. Confidence Intervals for the Difference of Two Means

In all cases we assume the data are:

$$y_{1,1}, y_{1,2}, \dots, y_{1,n_1} \sim N(\mu_1, \sigma_1^2), \text{ (population 1)}$$

$$y_{2,1}, y_{2,2}, \dots, y_{2,n_2} \sim N(\mu_2, \sigma_2^2), \text{ (population 2).}$$

We compute level L confidence intervals for $\mu_1 - \mu_2$.

Case 1: Paired Data

In this case, we take differences $d_i = y_{1,i} - y_{2,i}$. Then a confidence interval for the mean difference is also a confidence interval for $\mu_1 - \mu_2$.

Case 2: Independent Populations, Variances Assumed Equal

Assumption $\sigma_1^2 = \sigma_2^2$, and their value is unknown.

Formulas Estimate the population variance with the pooled variance estimator

$$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2},$$

where s_1^2 and s_2^2 are the sample standard deviations. The confidence interval is

$$\bar{y}_1 - \bar{y}_2 \pm t_{n_1+n_2-2, (1+L)/2} \sqrt{s_p^2 \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}.$$

Case 3: Independent Populations, Variances Not Assumed Equal

The confidence interval is

$$\bar{y}_1 - \bar{y}_2 \pm t_{\nu, (1+L)/2} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}},$$

where ν is taken as the largest integer less than or equal to $\left[\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right]^2 / \left[\frac{(s_1^2/n_1)^2}{n_1-1} + \frac{(s_2^2/n_2)^2}{n_2-1} \right]$.

IV. An Approximate Score Confidence Interval for the Difference of Two Proportions

Assumptions

1. We are interested in estimating $p_1 - p_2$, where p_1 is the proportion having a certain characteristic in population 1 and p_2 is the proportion having a certain characteristic in population 2.
2. y_1 is the number having the characteristic in a random sample of size n_1 taken from population 1 and y_2 is the number having the characteristic in a random sample of size n_2 taken from population 2.

Formula

A level L confidence interval for $p_1 - p_2$ is

$$\tilde{p}_1 - \tilde{p}_2 \pm z_{(1+L)/2} \sqrt{\frac{\tilde{p}_1(1-\tilde{p}_1)}{n_1} + \frac{\tilde{p}_2(1-\tilde{p}_2)}{n_2}},$$

where

$$\tilde{p}_1 = \frac{y_1 + 0.25z_{(1+L)/2}^2}{n_1 + 0.5z_{(1+L)/2}^2} \text{ and } \tilde{p}_2 = \frac{y_2 + 0.25z_{(1+L)/2}^2}{n_2 + 0.5z_{(1+L)/2}^2}.$$